



THE PREVALENCE OF COLOR VISION DISORDER (CVD) AND CONSANGUINITY IN POPULATION OF DISTRICT KARAK, KHYBER PAKHTUNKHWA (KP), PAKISTAN

Sawera Nisar¹, Muhammad Goher Zaman^{1*}, Faheem Ullah Khan², Hijab Nisa¹, Hira Maheen¹, Rubeena Gull¹, Shakir Ullah Khan¹

¹Department of Zoology, Government Post Graduate College Karak, KP, Pakistan

²Department of Zoology, University of Science and Technology Bannu, KP, Pakistan

ARTICLE INFO:

Keywords:

Color blindness, Ishihara test, City university test, Farnsworth D-15, consanguinity, X linked recessive, District Karak, KP, Pakistan.

Corresponding Author:

Muhammad Goher Zaman,
Department of Zoology,
Government Post Graduate
College Karak, KP, Pakistan
Email: goherzaman5@gmail.com

Article History:

Published on 03 August 2025

Authors contribution:

All authors made equal contributions to the conception and design of the project, data analysis and preparation of the manuscript.

ABSTRACT

Objectives: To observe the prevalence of color vision disorder (CVD) in district Karak, Khyber Pakhtunkhwa (KP) Pakistan from September 2023 to September 2024.

Methodology: During this field work, samples of 555 (300 male and 255 female) individuals were collected from surrounding locations, i.e. Karak city, Metha khel, Landiwa, Thth-e-Nasrati, Gurguri, Latamber, Banda Dawood Shah and Shagi Banda. The data were collected via a predesigned questionnaire printed in Urdu and English, and the history of consanguinity of parents was also recorded.

Results: We used Ishihara test, City university test (CUT) and Farnsworth D-15 to identify color blind people, revealed that 28 people (5.04%) were color blind. Out of these, 19(67.85%) were males and 9(32.14%) were females. Among these, 28 color blind individuals, 16(57.14%) were red-green color blind, 5 (17.855) were red color blind, 5 (17.85%) were green color blind and 2 (7.14%) were blue color blind. Furthermore, consanguinity of parents was recorded and interestingly all affected individual parents had consanguineous marriages. Two different types of pedigree analysis were recorded too. In the collected data, people aged between 18 to 60 years were present.

Conclusion: We found a significant prevalence of CVD in males (6.3%) than in females (3.5%). The study is significant in that it reported the prevalence of color blindness for the first time in Southern KP, Pakistan. This study provides the foundational understanding of CVD and its prevalence in district Karak and a framework for further research and prevention strategies regarding CVD.

INTRODUCTION

Worldwide, average color blindness prevalence is approximately 7 percent in males and 0.5 percent in females; however, its prevalence varies from one geographical region to another and from one race to another in different populations. Similarly, in India, prevalence of CVD is recorded as 3.89% in male and 0.18% in females¹. According to a comparative study conducted in Pakistan between medical and non-medical students, the total prevalence of color vision impairment was from 2.75 to 2.8%². Human eyes have two types of photoreceptor cells located on retina: the rod cells and the cone cells. The Rod cells are responsible for differentiation between black and white colors, or between day and night, whereas the cone cells are responsible for discrimination among different color³. Human eyes respond to light through a series of events called photo-transduction⁴.

Normally, blindness is caused by degenerative disorders of the retina called retinal dystrophies (RD) like disturbance in light sensation or response of photoreceptor cells to light, physical injury to eye or mutation in proteins responsible for normal vision. So, in this way, blindness is of two types. It may be complete blindness in which total vision loss occurs or it may be color blindness⁵⁻⁷. It may be caused due to disturbance in the cone cells, retina, photo perception or mutation in X chromosome⁸. Mostly, it is congenital and due to the missense mutation in genomic structure of L and M genes, located on long arm of the X chromosomes. Normally, the human eye can absorb three specific wavelength photons, i.e., red, green, and blue. Due to this absorbing capacity, normal human vision is trichromatic and is called trichromatic vision or trichromacy⁹. There are three types of color blindness: monochromatism, dichromatism, and achromatism. From molecular studies, it is concluded that color vision defects resulting from alteration, defects, absence, or malfunctioning of one photopigment result in dichromatism, two pigments absence or malfunctioning gives

rise to monochromatic, and total absence or malfunctioning of all three photopigments results in achromatism or total color vision loss¹⁰⁻¹².

Published summaries of all the data available before 1960s showed that the prevalence of color blindness in male population is about 8 per cent in European Caucasians, 5 per cent in Asians, 4% in Africans and less than 2 per cent in indigenous Americans, Australians, and Polynesians¹³. Similarly, the prevalence of CVD in Muslim countries also varies. Its frequency is recorded from 5.41% to 7.86% in different cities of Turkey¹⁴. In different cities of Saudi Arabia, the prevalence of CVD is recorded 5.85% in males and 0.75% in females in Riyadh, 1.77% in Makkah¹⁵. Similarly, in India Prevalence of CVD is recorded as 3.89% in males and 0.18% in females¹⁶. Mughal et al studied the prevalence of CVD in different countries and reported its prevalence 2.1% in 7,542 males and 0.2 % in 3,519 females in India. In Potohara, Western Nepal, 10 to 19 years age group children were studied and the prevalence of CVD was recorded as 3.8% in 18 boys but no female was reported to be colour blind¹⁷. In Iran, the recorded prevalence for the first time was 8.18% in 93 studied males and 4.93% in 53 males, while 0.43% in 4 studied females and 0.32% in 3 studied females¹⁸.


In Pakistan, the known percentages of CVD among males are 3.59%, 5.69%, 2.75%, 2.24%, 10.0%, 7.95 % and 3.1% and in females known percentages are 4.4%, 1.64%, and 1.39 %. It means that the prevalence of CVD varies from province to province in Pakistan and many factors contribute to this variation like consanguineous marriages, environmental factors, etc. Due to these factors, the frequency varies from province to province and even from district to district. In spite of all these problems, the prevalence of CVD reported from different regions of Pakistan in different published materials is 5.1% in Rawalpindi, 2.48% in Quetta, 3.1% in Southern Punjab¹⁹. In Karachi reported a total of 0.9% prevalence of CVD in 3437

individuals' sample. Among these CVD positive individuals, 1.45 were males and 0.4% were females. The prevalence of CVD in Peshawar was 2.91% in males and 0.83% in females among dental students at Sardar Begum Dental College Peshawar²⁰.

MATERIALS AND METHODS

Study Area

The study was conducted in different areas of District Karak, Khyber Pakhtunkhwa, Pakistan, from September 2023 to September 2024. The total area of Karak is 3372 sq km, and the total population is 815,878 (2023 Census). The majority of the people are very poor and have low economic status, and there is a lack of facilities regarding proper health and awareness about the disorder and its consanguineous transmission.



Study Design

The study was designed to know the prevalence of color vision disorder and its consanguinity factor in individuals of both sexes of all age groups in different regions in Karak including city and its surrounding areas such as Metha khel, Landiwa, Thath e Nasrati, Gurguri, Latamber, Banda Dawood Shah and Shagi Banda (Table 1).

Data Collection

The data was collected from different regions of district Karak after approval from the Ethical Review Committee of Government Postgraduate College, Karak (GPGC Karak). The sample size calculated was n=555. The study included all individuals' cases of vision disorder. A pre-designed questionnaire,

printed both in English and Urdu, was used to collect data. The history of consanguinity of parents was also recorded.

Table .1 Details of Data Collection

Serial No	Location	No of sample 2023-2024
1	Karak city	150
2	Metha khel	100
3	Tahth e Nasrati	70
4	Latamber	135
5	Landiwa	30
6	Gurguri	55
7	Banda Dawood Shah	12
8	Shagi Banda	3
Total		555

Screening

Total 38 different Ishihara plates were used for the detection of Red-Green monochromacy which contains various figures and lines²¹. while to find protanopia, deuteranopia and Tritanopia and making the diagnostic procedure more accurate and precise, City university test (CUT) and Farnsworth D-15 test were performed. CUT contained 10 plates to assess colour vision. Each plate features a central coloured dot surrounded by four peripheral dots, designed to test for various colour vision disorders, including those related to protan, deutan, and tritan deficiencies²². The Farnsworth D-15 test is a method for screening CVD by having the patient arrange coloured discs in a sequence, with a reference disc provided. It's a dichromatic test, meaning it's designed to distinguish between individuals with normal colour vision and those with mild to severe colour vision disorders. The test can help in identifying issues with red-green discrimination (deutan or protan) and blue-yellow discrimination (tritan)²³.

RESULTS

A total of 555 individuals were screened for color blindness both male and female. Before screening, a questionnaire was filled with all necessary information, including the personal profile (name, sex, age, profession, education, address, pedigree). In 555 individuals, 300 (54.05%) were males and 255 (45.94%) were females. Only 28 were found positive for CVD in both male and female individuals via diagnostic tests. Out of the 28 individuals, 19 (67.85%) were male and 9 (32.14%) were female. Among the 28 positive individuals, 16 (57.14%) were positive for red-green color blindness, 5 (17.85%) were positive for red color blindness. Whereas 5 (17.85%) were positive for green color blindness and 2 (7.14%) were positive for blue color vision disorder (Fig 1).

Fig 1. Total prevalence of CVD in population of District Karak







Figure 2. Location-wise prevalence of CVD in population of District Karak




(B)




(C)




(D)




(E)




(F)



(G)



(H)




Gender wise prevalence of CVD in population of District Karak


We noticed that number of CVD female ratio were more significant than male population as shown in Fig 3.

Fig 3. Gender wise Prevalence of CVD in population of District Karak

(A)



(B)







Age wise Prevalence of CVD in Male District Karak

Next, we examined the color blindness disorder on age basis. We found that 18-25 Protanopia 0%, Deuteranopia 1.11%, Tritanopia 0%, Red-Green monochromacy 4.44%. In age of 26-35, Protanopia 2.86%, Deuteranopia 1.43%, Tritanopia 0%, Red-Green monochromacy 5.71%.

The age of were 36-45 Protanopia 1.12%, Deuteranopia 2.25%, Tritanopia 0%, Red-Green monochromacy 2.25%. The age of 46-60 were Protanopia 1.96%, Deuteranopia

0%, Tritanopia 1.96%, Red-Green monochromacy 0% as shown in Fig.4.


Fig 4. Age wise Prevalence of CVD in Male District Karak




Age wise Prevalence of Color Blindness Disorder in Female District Karak

The age-wise female ratio was in of 18-25 Protanopia 0%, deuteranopia 0%, Tritanopia 0%, Red-Green monochromacy 0%. The age-wise female ratio was in of 26-35 Protanopia 0%, deuteranopia 0%, Tritanopia 1%, Red-Green monochromacy 0%. The age-wise female ratio was in of 36-45 Protanopia 4%, deuteranopia 4%, Tritanopia 0%, Red-Green monochromacy 4%. The age-wise female ratio was in of 46-60 Protanopia 0%, deuteranopia 0%, Tritanopia 0%, Red-Green monochromacy 5.71% as shown in Fig 5.


Fig 5. Age wise Prevalence of CVD in Female of District Karak




(B)



(C)



(D)



Consanguinity of the Disorder

In this study two different types of pedigrees were recorded. In one type pedigree, the female proband was identified with color blindness disorder who was married to a normal male, although they were not consanguineous related (Fig 6A). The result was two affected male individuals and four female carrier individuals were born. One of the carrier female individual was married to her close cousin who was normal. The resultant two male progenies were affected and one female was normal (Fig 6C).

Another type of pedigree was identified with male proband individual in our consanguinity related analysis.

Fig 6. Pedigree of One Female Proband Identified during Study





Fig 7. Pedigree of One Male Proband Identified during Study



When an affected male married to a normal woman, although they were not consanguineous related (Fig 7A), the result was a normal male and two carrier female individuals were born. One of the carrier's female as married to her close cousin who was affected then; two affected female individuals and one normal male individual were born (Fig 7C). Tritanopia is an autosomal dominant disorder so consanguinity is not important for its inheritance from parents to their offspring's. As in our study two individual's male and female were recorded, their parents were normal but they were affected with tritanopia (Table 2).

Table 2. CVD in relation to consanguinity in study Population

Consanguinity among parents of study individuals	Protanopia	Deuteranopia	Tritanopia	Red-Green monochromacy
Maternal Aunt Cousin (Khalazad)	0	2(M)	0	4(M)
Maternal Uncle Cousin (Mamonnzad)	1(M)	1(M)	1(M) (Mutated)	2(F)
Paternal Aunt Cousin (Phuphizad)	1(F)	1(M)	0	3(M)
Paternal Uncle Cousin (Chachazad)	2(M)	0	1(F) (Mutated)	2(M)
Second Cousin	0	0	0	3(F)
Distant Blood Relation	0	0	0	0
Same Sub-caste (Bradri)	1(M)	0	0	1(F)
Same Caste	0	1(F)	0	1(M)
Different Caste	0	0	0	0
Total	5	5	2	16

DISCUSSION

Under normal lighting circumstances, a failure or diminished ability to detect color variations is known as color blindness or color vision disorder (CVD). CVD is a predominant X linked recessive disorder having genes located on the long arm of the X chromosome at loci 28 (Xq28), so its prevalence in males would be more as compare to females²⁴. We examined 555 individuals among only 28 were positive. Among these positive, 19 were males and 9 were females. The previous studies revealed the same results of high prevalence rate of disorder in males 8% than females 0.4 %. Prevalence of CVD varies from area to area and caste to caste but its prevalence occurs mostly 3 to 8% in males and 0.3 to 0.5% in females and has been proved via many

literatures²⁵. In all forms of color blindness, red green color blindness is most common form, which is inherited as X linked recessive disorder²⁶.

Furthermore, we observed that the number of red color-blind individuals was similar to that of green color blind. Among 5 out of 28 (17.85%) were red color blind and 5 (17.85%) were green color blind and all affected individuals'

parents' marriages were consanguineous. Previous study has shown that protanopia was 1.94% and deuteranopia was 2.27%²⁷. It can be concluded that CVD prevalence may vary from one region to another and from one race to another race²⁸. Herein, the history of two families has been recorded in 8 locations of district Karak in which both consanguineous and inter-caste marriages were recorded. In 2019 dichromacy and color vision impairment study was done in district Chiniot Punjab, Pakistan where the affected individual parents' history was recorded in which 7 dichromatic individuals' parents were paternal uncle cousins (First cousin), 4 dichromatic parents were second cousins, 3 individual parents were maternal aunt cousins (second cousin), 1 individual parent was maternal uncle cousin (Mamuzad), 3 individual parents belonged to the same sub-caste (bradri), one parent was of the same caste while that of two individuals were of different castes. Their study and our current study have similarity in family history of affected individuals because in this study, recorded consanguinity rate 2 red green monochromats, 2 protanopia individual parents were paternal uncle cousins (Chachazad), 3 red green, 1 protanopic and 1 deuteranopic individual parents were paternal aunt cousins (Phuphizad), 4 red green color blind, and 2 deuteranopic individual parents were maternal aunt cousins (Khalazad), 3 red green defective individual parents were second cousins, 1 protanopic individual parent was of the same sub-caste (bradri) while that of one deuteranopic individual parent was of the same caste²⁹.

According to consanguinity analysis protanopia, deuteranopia and red-green monochromacy are X-linked recessive disorders while tritanopia is an autosomal chromosome number 7 linked dominant disorder³⁰. Although it is very rare even 1 in 13000 affected and can pass from parents to children with 50% probability. In our study 2 individuals one was male and another was female recorded. Although tritanopia is not

linked directly to consanguinity like other CVDs but consanguinity can increase the risk of certain autosomal dominant disorders³¹. The parents of tritanopic male and female individuals were maternal uncle-cousin and paternal uncle-cousin, they were normal but their children were tritanopic. This showed some type of autosomal mutation that affects male and female equally as recorded in this study and many other studies as well. As it is very rare, so it is not considered in most of the studies or is neglected. CVD being an X-linked recessive disorder is deeply related with gender and consanguinity, but it has no relation with age. In this study almost, all age groups from 18 to 60 were included and it showed that color blind has no relation with age³².

CONCLUSION

This study is significant for reporting the prevalence of color blindness for the first time in southern KP, Pakistan. The study also shows relation of CVD with consanguinity. This study provides basics, awareness and a pathway for further studies. Furthermore, it is also observed that CVD being X-linked recessive disorder is greatly related to consanguinity although, it may be onset disorder and may occur at any stage in life like other disease but mostly it is congenital. The findings reveal the prevalence of CVD as 6.3% in males and 3.5% in females of District Karak, KP, Pakistan. CVD although is not considered a serious disorder and even people are mostly unaware of its occurrence but sometime it becomes life threatening and becomes the rejection cause of people in many occupations like medical, arm forces etc. Till now no special treatment is discovered to treat CVD like other congenital disorders, and the patients remain unaware of their condition even throughout their life. This research provides a foundational understanding of CVD in district Karak KP Pakistan, providing a framework for further studies and prevention strategies.

REFERENCES

1. Warad C, Wani R, Singh R. Prevalence of Congenital Colour Vision Deficiency in

- Belagavi, Karnataka, India: A Hospital-based Cross-sectional Study. *J. Clin. Diagn. Res.* 2022 1;16(11).
2. Ahmed S, Chaudhary MM, Anwar A, Saad HT. A Comprehensive Cross-sectional Analysis of Prevalence of Congenital Color Vision Deficiency among Patients. *JLUMHS.* 2024 28;23(02):14650.
 3. Pandiyan VP, Maloney-Bertelli A, Kuchenbecker JA, Boyle KC, Ling T, Chen ZC, Park BH, Roorda A, Palanker D, Sabesan R. The optoretinogram reveals the primary steps of phototransduction in the living human eye. *Sci. Adv.* 2020 9;6(37): eabc1124.
 4. Kramer RH. Suppressing Retinal Remodeling to Mitigate Vision Loss in Photoreceptor Degenerative Disorders. *Annu. Rev. Vis. Sc.* 2023 15;9(1):131-53.
 5. B. Domènech E, Marfany G. The relevance of oxidative stress in the pathogenesis and therapy of retinal dystrophies. *Antioxid.* 2020 3;9(4):347.
 6. Rosa JG, Disner GR, Pinto FJ, Lima C, Lopes-Ferreira M. Revisiting retinal degeneration hallmarks: insights from molecular markers and therapy perspectives. *Int. J. Mol. Sc.* 2023 Aug 23;24(17):13079.
 7. Howorko L, Boedianto JM, Daniel B. The efficacy of stacked bar charts in supporting single-attribute and overall-attribute comparisons. *VISINF.* 2018 1;2(3):155-65.
 8. Boehm AE, Bosten J, MacLeod DI. Color discrimination in anomalous trichromacy: Experiment and theory. *Vision Res.* 2021 1; 188:85-95.
 9. Chandak NR, Daigavane SV, Sharma SR. Screening of colour vision deficiency in school children of Wardha District. *Indian J. Clin. Exp. Ophthalmol.* 2017 ;3(1):80-4.
 10. Gudeta TB, Asrat T. Prevalence and genotypic frequency of color vision defects among primary schoolchildren in Adama Town, Eastern Ethiopia. *BMC Pediatr.* 2024 22;24(1):72.
 11. Moreira H, Alvaro L, Lillo J. Color blindness and semantic knowledge: Cognition of color terms from elicited lists in dichromats and normal observers. *Color Res Appl.* 2024;49(5):420-32.
 12. Birch J. Worldwide prevalence of red-green color deficiency. *J. Opt. Soc. Am. A.* 2012 1;29(3):313-20.
 13. Allen K, Quinlan P, Andow J, Fischer E. What is it like to be colour-blind? A case study in experimental philosophy of experience. *MindLang.* 2022;37(5):8149.
 14. Althomali TA, Algarni FA, Alosaimi MA, Alharthi AA, Alharthi KA, Alotbi MF, Alharthi NS, Alsulimani AM, Aljuaid GM, Alsuwat AO. Prevalence of color blindness among secondary school students in Taif, Saudi Arabia. *Arch. Pharm. Pract.* 2022;13(3-2022):85-91.
 15. Kundu BK, Chakma B. Prevalence of colour vision defect in the indian population results from a preemployment screening centre of a tertiary care hospital. *J Contemp Med Res.* 2020;7(9):119-23.
 16. Niroula DR, Saha CG. The incidence of color blindness among some school children of Pokhara, Western Nepal. *Nepal Med Coll J.* 2010 1;12(1):48-50.
 17. Ugalahi MO, Fasina O, Ogun OA, Ajayi BG. Prevalence of congenital colour vision deficiency among secondary school students in Ibadan, South-West Nigeria. *Niger. Postgrad. Med. J.* 2016 1;23(2):93-6.
 18. Khan S, Rafique A, Khizer MA. Frequency of Colour Blindness amongst the Young Age Group in a Tertiary Care Eye Hospital. *Pak J Ophthalmol.* 2021 1;37(2):142-6.
 19. Ngente Z, Jadav B, Aafaque S, Patil A, Govindarajulu R, Muthusamy P, Babu JS, Swarnalatha C, Nayyar AS. Prevalence of

- color vision anomalies among dental professionals. *Middle East Afr. J. Ophthalmol.* 2021 1;28(3):169-73.
20. Ishihara S. The series of plates designed as tests for colour-blindness. Kanehara & Company; 1936.
 21. Karim KJ, Saleem MA. Prevalence of congenital red-green color vision defects among various ethnic groups of students in Erbil City. *Jordan J. Biol. Sci.* 2013 ;147(898):1-4.
 22. Hashemi H, Khabazkhoob M, Pakzad R, Yekta A, Heravian J, Nabovati P, Ostadimoghaddam H. The prevalence of color vision deficiency in the northeast of Iran. *J. Curr. Ophthalmol.* 2019 1;31(1):80-5.
 23. Janáky M, Braunitzer G. Syndromic Retinitis Pigmentosa: A Narrative Review. *J. Vis.* 2025 20;9(1):7.
 24. Moudgil T, Arora R, Kaur K. Color vision impairment in school children. *Highlights on Med. Res.* 2021 26;15:9-17.
 25. Fareed M, Anwar MA, Afzal M. Prevalence and gene frequency of color vision impairments among children of of six populations from North Indian region. *G&D.* 2015 1;2(2):211-8.
 26. El Moussawi Z, Boueiri M, Al-Haddad C. Gene therapy in color vision deficiency: a review. *Int. Ophthalmol.* 2021 ;41(5):1917-11827.
 27. Mitiku RG, Tolera BS, Tolesa ZG. Prevalence and allele frequency of congenital colour vision deficiency (CCVD) among students at Hawassa University, J. Egypt. Public Health Assoc. 2020 20;95(1):10.
 28. Zar MS, Akhtar MS, Haris AR, Aslamkhan M. Colour Vision Deficiency and Consanguinity in Pakistani Pukhtoon Population. *Adv. life sci.* 2020 2;7(4):237-9.
 29. Kim YJ, Packer O, Dacey DM. A circuit motif for color in the human foveal retina. *Proc Natl Acad Sci USA.* 2024 3;121(36):e2405138121.
 30. Stockman A, Sharpe LT. Human cone spectral sensitivities and color vision deficiencies. Totowa, NJ: Humana Press. 2008(pp.307327).
 31. Baraas RC, Hagen LA, Dees EW, Neitz M. Substitution of isoleucine for threonine at position 190 of S-opsin causes S-cone function abnormalities. *Vis. Res.* 2012 1;73:19
 32. Faiz S, Zehra S, Qureshi F, Mumtaz S. Colour blindness among male professional drivers: a cross-sectional study in Karachi. *Journal of Bahria University Medical and Dental College. JBUMDC.* 2019;9(2):124.